

Maricopa Family Dentistry & Orthodontics

Surgical Implant

Post- Treatment Instructions

For proper healing, it is very important that your intake of food and water be adequate. Swelling or discomfort may occur during the first 24-48 hours following placement of your Implant. Please follow these instructions:

- Avoid smoking and alcoholic beverages until healing is well established, as these activities will contribute to post-operative complications.
- Avoid excessive activities for the next 24 hours
- You may use ice packs as needed to control swelling for the first 24 hours.
- An antibiotic and anti-inflammatory may be prescribed to reduce swelling and prevent infection around surgical site; please take as prescribed.
- Use your Chlorhexidine mouth rinse and continue your usual oral hygiene program with care near the treatment area(s) and/ or suture(s).
- If swelling persists don't hesitate to call our office with any concerns.

If a pain medication has been prescribed by the doctor avoid driving and operating heavy machinery while taking the prescribed pain medication.

REMEMBER - A clean mouth heals faster!

If you have any questions or concerns do not hesitate to call our office at the number below. If you need immediate attention please call Dr. Pope at 480-332-6111.

◆ Doctor Jared Pope DDS ◆

◆ 44480 W. Honeycutt Ave. Suite 110 ◆ Maricopa, AZ. 85239 ◆

◆ Main 520-568-9100 ◆ Fax 520-568-9190 ◆

Maricopa Family Dentistry & Orthodontics

Surgical Implant

Post- Treatment Instructions

For proper healing, it is very important that your intake of food and water be adequate. Swelling or discomfort may occur during the first 24-48 hours following placement of your Implant. Please follow these instructions:

- Avoid smoking and alcoholic beverages until healing is well established, as these activities will contribute to post-operative complications.
- Avoid excessive activities for the next 24 hours
- You may use ice packs as needed to control swelling for the first 24 hours.
- An antibiotic and anti-inflammatory may be prescribed to reduce swelling and prevent infection around surgical site; please take as prescribed.
- Use your Chlorhexidine mouth rinse and continue your usual oral hygiene program with care near the treatment area(s) and/ or suture(s).
- If swelling persists don't hesitate to call our office with any concerns.

If a pain medication has been prescribed by the doctor avoid driving and operating heavy machinery while taking the prescribed pain medication.

REMEMBER - A clean mouth heals faster!

If you have any questions or concerns do not hesitate to call our office at the number below. If you need immediate attention please call Dr. Pope at 480-332-6111.

◆ Doctor Jared Pope DDS ◆

◆ 44480 W. Honeycutt Ave. Suite 110 ◆ Maricopa, AZ. 85239 ◆

◆ Main 520-568-9100 ◆ Fax 520-568-9190